



SOYBEAN DISEASE MANAGEMENT

Fungicide Efficacy for Control of Soybean Seedling Diseases

The members North Central Regional Committee on Soybean Diseases (NCERA-137)

have developed the following ratings for how well fungicide seed treatments control seedling diseases of soybeans in the United States. Efficacy ratings for each fungicide active ingredient listed in the table were determined by field-testing the materials over multiple years and locations by the members of this group, and include ratings summarized from national fungicide trials published in Plant Disease Management Reports (and formerly Fungicide and Nematicide Tests) by the American Phytopathological Society at http:// www.apsnet.org. Each rating is based on the fungicide's level of disease control, and does not necessarily reflect efficacy of fungicide active ingredient combinations and/or yield increases obtained from applying the active ingredient.

The list includes the most widely marketed products available. It is not intended to be a list of all labeled active ingredients and products. Additional active ingredients may be available, but have not been evaluated in a manner allowing a rating. Products listed are the most common products available as of the release date of the table; all available products may not be listed. Additional active ingredients may be included in some products for insect and nematode control, however; only active ingredients for pathogen control are listed and rated.

Many active ingredients and their products have specific use restrictions. Read and follow all use restrictions before applying any fungicide to seed, or before handling any fungicide-treated seed. This information is provided only as a guide. It is the applicator's and users legal responsibility to read and follow all current label directions. Reference in this publication to any specific commercial product, process, or service, or the use of any trade, firm, or corporation name is for general informational purposes only and does not constitute an endorsement, recommendation, or certification of any kind by members of the group, or by the North Central Soybean Research Program. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

Please note: Efficacy ratings may be dependent on the rate of the fungicide product on seed. A number of different species of *Pythium* and *Fusarium* impact seed and seedling health in soybean. Therefore, wide ranges in efficacy may be observed in fungicide active ingredients listed in the table. This is why several fungicide active ingredients are combined in seed treatments to provide protection to a broader spectrum of pathogens. Contact your local Extension plant pathologist for recommended fungicide product rate information for your area.



Seedling damping off from Pythium seedling blight and root rot Image: Martin Chilvers

Find Out More

The Crop Protection Network (CPN) is a multistate and international collaboration of university and provincial extension specialists, and public and private professionals who provide unbiased, research-based information to farmers and agricultural personnel. Our goal is to communicate relevant information that will help professionals identify and manage field crop diseases.

Find more crop disease resources at

CropProtectionNetwork.org

This publication was developed by members of NCERA-137. It was compiled by Kiersten Wise, University of Kentucky.

The information in this publication is only a guide, and the authors assume no liability for practices

implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer. ©**2022** by the Crop Protection Network. All rights reserved.



We Are Extension



ETWORK inversities

Fungicide Efficacy for Control of Soybean Seedling Diseases (02/2022)

Efficacy categories:

P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; NL=Not Labeled for use against this disease; NR=Not Recommended; U=Unknown efficacy or insufficient data to rank product

Fungicide active ingredient	Pythium spp. ¹	Phytophthora	Rhizoctonia spp.	Fusarium spp. ^{1,3}	Sudden death syndrome (SDS) Fusarium virguliforme	Phomopsis spp.
Azoxystrobin	P-G	NS	VG	F-G	NR	Р
Carboxin	U	U	G	U	NR	U
Ethaboxam	E	E	NR	NR	NR	NR
Fludioxonil	NR	NR	G	F-VG	NR	G
Fluopyram	NR	NR	NR	NR	VG	NR
Fluxapyroxad	U	U	E	G	NR	G
Ipconazole	Р	NR	F-G	F-E	NR	G
Mefenoxam	E ²	E	NR	NR	NR	NR
Metalaxyl	E ²	E	NR	NR	NR	NR
Oxathiapiprolin	P-G	E	NR	NR	NR	NR
PCNB	NR	NR	G	U	NR	G
Penflufen	NR	NR	G	G	NR	G
Prothioconazole	NR	NR	G	G	NR	G
Pydiflumetofen	NL	NL	NL	NL	VG	NL
Pyraclostrobin	P-G	NR	F-G	F	NR	G
Sedaxane	NR	NR	E	NS	NR	G
Thiabendazole	NR	NR	NL	NL	Р	G
Trifloxystrobin	Р	Р	F-E	F-G	NR	P-F
Extract of <i>Chenopodium</i> <i>quinoa</i> saponins ⁴	U	U	U	U	P-F	U

¹ Products may vary in efficacy against different *Fusarium* and *Pythium* species. ² Areas with mefenoxam or metalaxyl insensitive populations may see less efficacy with these products. ³ Listed seed treatments do not have efficacy against *Fusarium virguliforme*, causal agent of sudden death syndrome. ⁴ This product is a biological seed treatment that is registered and approved for organic use.



The CPN would like to thank the United Soybean Board for their support of this publication.

Common Fungicide Trade Names and Active Ingredients (02/2022)

Product trade name	Active ingredient(s)			
Acceleron	DX-612 Fluxapyroxad, DX-309 Metalaxyl, DX-109 Pyraclostrobin			
Allegiance FL	Metalaxyl			
Allegiance LS	Metalaxyl			
Apron XL LS	Mefenoxam			
ApronMaxx RFC				
ApronMaxx RTA				
CruiserMaxx	Fludioxonil, Mefenoxam			
CruiserMaxx Advanced or Cruiser Maxx Plus				
CruiserMaxx Vibrance or Vibrance Trio	Fludioxonil, Mefenoxam, Sedaxane			
Dynasty	Azoxystrobin			
EverGol Energy SB	Metalaxyl, Penflufen, Prothioconazole			
Heads Up	Extract of Chenopodium quinoa saponins			
ILEVO	Fluopyram			
Inovate Pro	Ipconazole, Metalaxyl			
Intego	Ethaboxam			
Lumisena	Oxathiopiprolin, Metalaxyl			
Maxim 4FS	Fludioxonil			
Mertect 340 F	Thiabendazole			
Prevail	Carboxin, Metalaxyl, PCNB			
Saltro	Pydiflumetofen			
Trilex 2000	Metalaxyl, Trifloxystrobin			
Vibrance	Sedaxane			
Warden CX	Fludioxonil, Mefenoxam, Sedaxane			
Warden RTA	Fludioxonil, Mefenoxam			