



CPN-1019-W | Management of Soybean Foliar Diseases

Fungicide Efficacy for Control of Soybean Foliar Diseases

The North Central Regional Committee on Soybean Diseases (NCERA-137) has developed the following information on foliar fungicide efficacy for control of major foliar soybean diseases in the United States. Efficacy ratings for each fungicide listed in the table were determined by field-testing the materials over multiple years and locations by the members of the committee. Efficacy ratings are based upon level of disease control achieved by product and are not necessarily reflective of yield increases obtained from product application. Efficacy depends upon proper application timing, rate, and application method to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in the table, unless otherwise noted. For application timing and use considerations, please contact your local cooperative extension service. Table includes systemic fungicides available that have been tested over multiple years and locations. The table is not intended to be a list of all labeled products¹.



Find Out More

The Crop Protection Network (CPN) is a multi-state and international collaboration of university and provincial extension specialists, and public and private professionals who provide unbiased, research-based information to farmers and agricultural personnel. Our goal is to communicate relevant information that will help professionals identify and manage field crop diseases.

Find more crop disease resources at CropProtectionNetwork.org.



This publication was developed by members of NCERA-137. It was compiled by Kiersten Wise and Luke Warner, University of Kentucky.

The information in this publication is only a guide, and the authors assume no liability for practices implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](https://www.usda.gov/program-discrimination-complaint) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

©2026 by the Crop Protection Network. All rights reserved.

Fungicide Efficacy for Control of Soybean Foliar Diseases Table (02/2026)

Efficacy categories: P = Poor; F = Fair; G = Good; VG = Very Good; E = Excellent; NL = Not labeled for use against this disease; U = Labeled, but unknown efficacy or insufficient data to rank product.

Fungicide mode of action groups:

Group 11 QoI Strobilurins | Group 1 MBC Thiophanates | Group 29 2,6-Dinitro-anilines | Group 3 DMI Triazoles | Group 7 SDHI Carboxamides | Group BM-01 Plant Extracts

Indicates product with mixed fungicide classes

	Active ingredient(%) ¹	Product/Trade name	Rate/A (fl oz)	Commonly tested rate/A (fl/oz)	PHI	Aerial web blight ²	Brown spot ³	Cercospora leaf blight ⁴	Frogeye leaf spot ⁵	Diaporthe (Pod and stem blight)	Soybean rust	Target spot	White mold ⁶
11	Azoxystrobin 22.9%	Quadris 2.08 SC, multiple generics	6.0-15.5	6	14	VG	P-G	P	P	U	G-VG	NL	NL
	Fluoxastrobin 40.3%	Aftershock 480 SC, Evito 480 SC	2.0-5.7	4	30	NL	P-G	P	P	NL	U	NL	NL
	Picoxystrobin 22.5%	Approach 2.08 SC	6.0-12.0	9	14	VG	P-G	P	P	U	G	U	G ⁷
	Pyraclostrobin 23.6%	Headline 2.09 EC/SC	6.0-12.0	12	21	NL	P-G	P	P	U	VG	NL	NL
3	Cyproconazole 8.9%	Alto 100SL	2.75-5.5	NA	14	U	VG	F	NL	U	VG	NL	NL
	Flutriafol 11.8%	Topguard 1.04 SC	7.0-14.0	7	21	NL	VG	P-G	G-VG	NL	VG-E	P	NL
	Propiconazole 41.8%	Tilt 3.6 EC, multiple generics	4.0-6.0	4	R5	P	G	NL	F	NL	VG	NL	NL
	Prothioconazole 41.0%	Proline 490 SC ⁸	2.5-5.0	5	21	NL	NL	NL	G-VG	NL	VG	NL	F
	Tetraconazole 20.5%	Domark 230 ME, multiple generics	4.0-5.0	5	R5	NL	VG	P-G	F-G	NL	VG-E	NL	F
1	Thiophanate-methyl 70%	Topsin-M, multiple generics	10.0-20.0	20	21	U	U	NL	G-VG	U	NL	NL	F
29	Fluazinam 40.0%	Omega 500 DF	12.0-16.0	16	30	NL	NL	NL	NL	NL	NL	NL	G
7	Boscalid 70%	Endura 0.7 DF	3.5-11.0	8	21	NL	NL	U	P	NL	NL	NL	VG
	Inpyrflumax 31.25%	Excalia 2.84 SC	2	2	R5	E	NL	NL	NL	NL	U	NL	NL
11	Azoxystrobin 25.30%	Topguard EQ 4.29 SC	5.0-8.0	5	21	VG	VG	U	G-VG	U	E	P	U
3	Flutriafol 18.63%												
11	Azoxystrobin 18.2%	Quadris Top 2.72 SC	8.0-14.0	8	14	U	G-VG	P-G	G-VG	F-G	VG	NL	NL
3	Difenoconazole 11.4%												
11	Azoxystrobin 19.8%	Quadris Top SBX 3.76 SC	7.0-7.5	7	14	VG	G-VG	P-G	G-VG	F-G	VG	NL	NL
3	Difenoconazole 19.8%												
11	Azoxystrobin 7.0%	Quilt 1.66 SC, multiple generics	14.0-20.5	NA	NA	U	G	F	F	U	VG	NL	NL
3	Propiconazole 11.7%												
11	Azoxystrobin 13.5%	Quilt Xcel 2.2 SE, multiple generics	10.5-21.0	10.5	R6	E	G	F	F	U	VG	NL	NL
3	Propiconazole 11.7%												
7	Benzoindiflupyr 2.9%	Trivapro 2.21 SE	13.7-20.7	13.7	14	E	G-VG	P-G	F-G	G	VG-E	U	NL
11	Azoxystrobin 10.5%												
3	Propiconazole 11.9%												
3	Cyproconazole 7.17%	Approach Prima 2.34 SC	5.0-6.8	6.8	30	VG	G	P-G	F-G	U	VG-E	F-G	NL
11	Picoxystrobin 17.94%												
7	Fluopyram 17.4%	Propulse 3.34 SC	6.0-10.2	8	21	NL	U	NL	U	U	U	NL	G
3	Prothioconazole 17.4%												
7	Bixafen 15.55%	Lucento 4.17 SC	3.0-5.5	5	21	NL	VG	F-G	G-VG	NL	VG-E	F-G	U
3	Flutriafol 26.47%												
11	Fluoxastrobin 14.84%	Fortix 3.22 SC, Preemptor SC	4.0-6.0	5	30	U	G-VG	P-G	G-VG	U	U	NL	U
3	Flutriafol 19.3%												

¹Multiple fungicides are labeled for soybean rust only, powdery mildew, and Alternaria leaf spot, including tebuconazole (multiple products) and myclobutanil (Laredo). Contact fungicides such as chlorothalonil may also be labeled for use. ²In areas where QoI-fungicide resistant isolates of the aerial blight pathogen are present, QoI fungicides may result in poor disease control. ³In areas where QoI-fungicide resistant isolates of the brown spot pathogen are present, QoI fungicides may result in poor disease control. ⁴Cercospora leaf blight efficacy relies on accurate application timing, and standard R3 application timings may not provide adequate disease control. Fungicide efficacy may improve with earlier or later applications; however, efficacy has been inconsistent with some products. Fungicides with a solo or mixed QoI or MBC mode of action may not be effective in areas where QoI or MBC resistance has been detected in the fungal population that causes Cercospora leaf blight. ⁵In areas where QoI-fungicide resistant isolates of the frogeye leaf spot pathogen are not present, QoI fungicides may be more effective than indicated in this table. ⁶White mold efficacy is based on R1-R2 application timing, and lower efficacy is obtained at R3 or later application timings, or if disease symptoms are already present at the time of application. ⁷Rating is based on two applications of a 9 fl oz/A rate of Approach at R1 and R3. ⁸Proline has a supplemental label (2ee) for white mold in NY. ⁹Stratego YLD has a supplemental label (2ee) for white mold on soybean only in IL, IN, IA, MI, MN, NE, ND, OH, SD, WI.

Fungicide Efficacy for Control of Soybean Foliar Diseases Table (02/2026)

Efficacy categories: P = Poor; F = Fair; G = Good; VG = Very Good; E = Excellent; NL = Not labeled for use against this disease; U = Labeled, but unknown efficacy or insufficient data to rank product.

Fungicide mode of action groups:

Group 11 QoI Strobilurins | Group 1 MBC Thiophanates | Group 29 2,6-Dinitro-anilines | Group 3 DMI Triazoles | Group 7 SDHI Carboxamides | Group BM-01 Plant Extracts

	Active ingredient (%) ¹	Product/Trade name	Rate/A (fl oz)	Commonly tested rate/A (fl/oz)	PHI	Aerial web blight ²	Brown spot ³	Cercospora leaf blight ⁴	Frogeye leaf spot ⁵	Diaporthe (Pod and stem blight)	Soybean rust	Target spot	White mold ⁶
11	Trifloxystrobin 13.7%	Delaro 325 SC	8.0-11.0	8	21	VG	VG	U	G-VG	U	U	NL	F
3	Prothioconazole 16.0%												
7	Fluopyram 10.9%	Delaro Complete 3.83 SC	8.0-11.0	8	21	U	VG	U	G-VG	U	U	NL	F
11	Trifloxystrobin 13.1%												
3	Prothioconazole 14.9%												
7	Pydiflumetofen 6.9%	Miravis Top 1.67 SC	13.7	13.7	14	VG	VG	F-G	G-VG	G	NL	F-G	U
3	Difenoconazole 11.5%												
7	Pydiflumetofen 7.0%	Miravis Neo 2.5 SC	13.7-20.8	13.7	14	U	U	U	G-VG	U	U	U	P
11	Azoxystrobin 9.3%												
3	Propiconazole 11.6%												
11	Pyraclostrobin 28.58%	Priaxor 4.17 SC	4.0-8.0	4	21	NL	G-VG	P-G	P-F	U	VG-E	F-G	P
7	Fluxapyroxad 14.33%												
11	Trifloxystrobin 32.3%	Stratego YLD 4.18 SC ⁹	4.0-4.65	4	21	VG	G	F	F-G	U	VG	P	NL ⁹
3	Prothioconazole 10.8%												
11	Azoxystrobin 9.35%	Affiance 1.5 SC	10.0-14.0	10.0-14.0	14	U	VG	F	F-G	U	U	NL	U
3	Tetraconazole 7.48%												
11	Fluoxastrobin 17.76%	Zolera FX 3.34 SC	4.4-6.8	5	30	U	U	U	F-G	U	U	NL	U
3	Tetraconazole 17.76%												
1	Thiophanate-methyl 21.27%	Acropolis 2.37 SC	20.0-23.0	23	R5	NL	U	U	G-VG	U	NL	NL	U
3	Tetraconazole 4.20%												
7	Fluxapyroxad 7.74%	Revytek 3.33 SC	8.0-15.0	8	21	VG	VG	F-VG	G-VG	U	VG-E	F-VG	P
11	Pyraclostrobin 15.49%												
3	Mefentrifluconazole 11.61%												
11	Pyraclostrobin 17.56%	Veltyma 3.34 SC	7.0-10.0	7	21	U	U	U	G-VG	U	U	NL	NL
3	Mefentrifluconazole 17.56%												
BM-01	Tea Tree Oil 20.4%	Regev HBX 3.34 EC	4.0-8.5	8.5	14	NL	U	U	G-VG	U	U	NL	U
3	Difenoconazole 20.4%												
1	Thiophanate-methyl 37.50%	Frogghom	20	20	21	U	U	NL	G-VG	U	U	NL	U
3	Tebuconazole 7.50%	Overrule											
3	Mefentrifluconazole 26.04%	Revylok 3.33 SC	4.5-6.5	5.5	21	U	U	U	G-VG	U	U	U	U
7	Fluxapyroxad 8.68%												
11	Picoxystrobin 17.05%	Viatitude 2.09 SC	8.0-16.0	NA	36	U	U	U	G	U	U	U	F
3	Prothioconazole 5.68%												
7	Boscalid 18.00%	Zorina	18.5-20	NA	21	U	U	U	U	U	U	NL	U
3	Mefentrifluconazole 9%												

¹Multiple fungicides are labeled for soybean rust only, powdery mildew, and Alternaria leaf spot, including tebuconazole (multiple products) and myclobutanil (Laredo). Contact fungicides such as chlorothalonil may also be labeled for use. ²In areas where QoI-fungicide resistant isolates of the aerial blight pathogen are present, QoI fungicides may result in poor disease control. ³In areas where QoI-fungicide resistant isolates of the brown spot pathogen are present, QoI fungicides may result in poor disease control. ⁴Cercospora leaf blight efficacy relies on accurate application timing, and standard R3 application timings may not provide adequate disease control. Fungicide efficacy may improve with earlier or later applications; however, efficacy has been inconsistent with some products. Fungicides with a solo or mixed QoI or MBC mode of action may not be effective in areas where QoI or MBC resistance has been detected in the fungal population that causes Cercospora leaf blight. ⁵In areas where QoI-fungicide resistant isolates of the frogeye leaf spot pathogen are not present, QoI fungicides may be more effective than indicated in this table. ⁶White mold efficacy is based on R1-R2 application timing, and lower efficacy is obtained at R3 or later application timings, or if disease symptoms are already present at the time of application. ⁷Rating is based on two applications of a 9 fl oz/A rate of Approach at R1 and R3. ⁸Proline has a supplemental label (2ee) for white mold in NY. ⁹Stratego YLD has a supplemental label (2ee) for white mold on soybean only in IL, IN, IA, MI, MN, NE, ND, OH, SD, WI.

Many products have specific use restrictions about the amount of active ingredient that can be applied within a period of time or the amount of sequential applications that can occur. Please read and follow all specific use restrictions prior to fungicide use and follow all harvest restrictions provided on the label. This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Persons using such products assume responsibility for their use in accordance with current directions of the manufacturer. Members or participants in the NCERA-137 group assume no liability resulting from the use of these products.

This article is published by the Crop Protection Network with funding provided by U.S. soybean farmers through the United Soybean Board, the United States Department of Agriculture - National Institute of Food and Agriculture, and the Grain Farmers of Ontario through the Canadian Agricultural Partnership (CAP), a federal-provincial territorial initiative.