



**CROP
PROTECTION
NETWORK**

CPN-7006 | Cotton Disease Management

Nematicide Efficacy for Control of Cotton Nematodes

The members of the Cotton Nematologists Working Group (CNWG) have developed the following ratings for the effectiveness of nematicides in controlling nematodes of cotton in the United States. Efficacy ratings for each nematicide listed were determined through field experiments conducted over multiple years and locations by members of the group. Ratings are based on control of nematode root galling, and/or reproduction, and do not necessarily reflect yield protection. Numerous factors influence nematicide performance in the field, such as soil moisture at and for several weeks after application, soil texture in both the upper 10 to 12 inches and deeper layers of the profile, and nematode population density. In general, nematicides applied in-furrow at-planting provide more consistent nematode control than seed treatments. Seed treatment nematicides are suggested in fields with low nematode densities, while in-furrow nematicides are suggested for fields with low to moderate nematode densities. Furthermore, host plant resistance is the preferred control measure to

utilize when nematode densities are high; however, host plant resistance is currently only available for the southern root-knot nematode and the reniform nematode in cotton. Host plant resistance is usually more cost-effective than using a nematicide. All nematicides are limited in their ability to prevent yield losses when cotton nematode densities are high. Nematicides can be cost effective for controlling lance or sting nematodes on root-knot or reniform nematode resistant varieties.



Find Out More

The Crop Protection Network (CPN) is a multi-state and international collaboration of university and provincial extension specialists, and public and private professionals who provide unbiased, research-based information to farmers and agricultural personnel. Our goal is to communicate relevant information that will help professionals identify and manage field crop diseases. Find more crop disease resources at CropProtectionNetwork.org.



**CROP
PROTECTION
NETWORK**

We Are Extension

This publication was developed by members of Cotton Nematologists Working Group and compiled by Travis Faske, University of Arkansas.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](https://www.usda.gov/program-discrimination-complaint) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

©2026 by the Crop Protection Network. All rights reserved.

Nematicide Efficacy for Control of Cotton Nematodes Table (04/2026)

The list includes the most common, commercially available products marketed as nematicides in cotton. Other nematicides may be commercially available, but there is not sufficient data to support a rating. Furthermore, all products are tested following the application methods listed on the most current US EPA label. Many of these products have restrictions outlined on the label, which are not listed here, so review the label before applying these products in cotton fields. Some products are sold as a premix with an insecticide or are also marketed as a fungicide; however, the ratings listed are an assessment of the product as a nematicide only.

Efficacy categories: P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; NL = Not Labeled; NR=Not Recommended; U = Unknown efficacy or insufficient data to rank product

Active ingredient (%)	Formulation ¹	Trade name	Rates	Commonly tested rates	Southern root-knot nematode	Reniform nematode	Lance nematode ²	Sting nematode
Abamectin 46.3%	Seed treatment	Avicta 500FS	0.15 mg ai/seed	N. A.	P-F	P-F	P	P
Thiodicarb 24% + imidacloprid 24%	Seed treatment	Aeris 5FS	0.75 mg ai/seed	N. A.	P	P	P	P
Clothianidin 40.3% + <i>Bacillus firmus</i> I-1582 8.1%	Seed treatment	Poncho/Votivo 5FS	0.424 mg ai/seed	N. A.	P	P	P	P
<i>Burkholderia</i> spp. A396	Seed treatment	BioST Nematicide 100	7.0 fl oz/cwt	N. A.	P-F	P-F	P-F	P-F
<i>Bacillus amyloliquefaciens</i> MBI 600 1% <i>cis-Jasmone</i> 0.88%	Seed treatment	Trunemco Cotton	0.004 - 0.21 mg ai/seed	0.21	P	P	P	P
Fluopyram 49.02%	Seed treatment	Copeo 600FS	0.2-0.3 mg ai/seed	0.2	F	F	F	P-F
Fluopyram 29.5% + <i>Bacillus firmus</i> I-1582 11.5%	Seed treatment	Acceleron N-314 4.17FS	0.24 - 0.49 mg ai/seed	0.3	U	U	U	U
<i>Bacillus licheniformis</i> FMCH001 + <i>Bacillus subtilis</i> FMCH002	Liquid - IF	Zironar	6.0-12.0 fl oz/ac	6.0	P	P	P	P
Fluopyram 41.5%	Liquid - IF	Velum 4.16SC	5.0-6.8 fl oz/A	6.0	G	F	F	F
Abamectin 8%	Liquid - IF	Averland 0.7FC	3.5 fl oz/A	3.0	F	F	P-F	P-F
Oxamyl 42% ³	Liquid - Foliar	Vydate C-LV	8.5-17.0 fl oz/A	17.0	P	P	P	P
Aldicarb 15%	Granular -IF	AgLogic 15GG	3.5-7.0 lb/A	5.0	G-VG	G-VG	F-VG	F-G
1,3-dichloropropene 97.5%	Liquid - Fumigant	Telone II 9.85L	3.0-6.0 gal/A	3.0	VG	VG	VG	VG

¹ Formulations: Seed treatment = applied on seed coat; Liquid - IF - liquid product applied in-furrow with 5- 6 gal/ac water carrier; Liquid-Foliar = liquid product applied broadcast; Liquid-fumigant = soil applied nematicides that are injected into the soil preplant; Granular-IF are granular materials applied in-furrow.

² Includes both *Hoplolaimus columbus* and *H. galeatus*

³ Vydate C-LV is applied with a granular- or liquid-IF nematicide or seed treatment. Foliar applications range from broadcast of 17 fl oz/ac at 1st to 7th true-leaf growth stage or banded application. See label for specific rates for these foliar applications and through drip irrigation. Rating based on foliar application only.

This information is provided only as a guide. It is the applicator's legal responsibility to read and follow all current label directions. Reference in this publication to any specific commercial product is for general information only and does not constitute an endorsement or recommendation by the CNWG. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer. Members of the CNWG assume no liability resulting from the use of these products.